

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/274898722>

Evaluation of Moringa Oliefera (Lam) (Moringaceae) Seed oil for Laval control of Aedes Aegypti L. (Diptera: Culicidae).

Article · November 2014

CITATION

1

READS

164

9 authors, including:



Nj Okonkwo

Nnamdi Azikiwe University, Awka

22 PUBLICATIONS 57 CITATIONS

[SEE PROFILE](#)



Edith Nwankwo

Nnamdi Azikiwe University, Awka

30 PUBLICATIONS 25 CITATIONS

[SEE PROFILE](#)



Ibeabuchi Uko

Nnamdi Azikiwe University, Awka

10 PUBLICATIONS 1 CITATION

[SEE PROFILE](#)



Chikaodili Ukonze

Nnamdi Azikiwe University, Awka

14 PUBLICATIONS 11 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Seasonal distribution and micro-climatic factors influencing the abundance of malaria vectors in the South-East Nigeria [View project](#)



Environmental pollution in relation to e waste [View project](#)

THE INTERNATIONAL JOURNAL OF SCIENCE & TECHNOLEDGE

Evaluation of *Moringa Oleifera* (Lam) (Moringaceae) Seed Oil For Larval Control of *Aedes Aegypti* L. (Diptera: Culicidae)

Okonkwo, Nnaemeka J

Department of Crop Science and Horticulture, Nnamdi Azikiwe University, Awka

Nwankwo, Edith N

Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka

Uko Ibeabuchi

Department of Crop Science and Horticulture, Nnamdi Azikiwe University, Awka

Okafor, Ebube G

Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka

Ukonze, Blessing C

Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka

Ogbonna, Confidence U

Department of Crop Science, University of Ghana, Legon, Accra, Ghana

Egbuche, Chukwudi M

Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka

Chikezie, Friday M

National Arbovirus and Vectors Research Centre, Enugu

Obiechina, Izuchukwu O

Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka

Abstract:

Comparative bioassays were carried out with 1st and 4th instar larvae of Aedes aegypti L. (Diptera: Culicidae) using Moringa oleifera seed oil. This was to investigate whether the oil extract would show lethal effects on the mosquito larvae and to further establish at which point during the larval growth the oil would be effective. The larvae were exposed at ambient laboratory temperatures of $28 \pm 2^{\circ}\text{C}$, $80 \pm 5\%$ r.h and photoperiod of 12:12 light and dark hours. Seven dosages of 200, 100, 50, 25, 12.5, 6.25 and 3.125 μl of oil dissolved in acetone per millilitre of water and control treatments with ordinary acetone were included in the trials. These were arranged on the laboratory bench in a Completely Randomised manner. Larval mortality was recorded at 3, 6, 9, and 12 hours post treatment. Mortality data were subjected to analysis of variance and log-probit regression, accordingly. Results showed that significant ($P < 0.05$) dosage-related mortality responses were noted for both instars. Thus, Moringa oil showed significant ($P < 0.05$) toxic action against all larval instars of Ae. aegypti tested. Higher concentrations of the oil resulted in higher kill of the larvae within 12 hours, while sublethal dosages gave low mortality rates. The LD_{50} values determined were 208.35 $\mu\text{l/ml}$ and 62.94 $\mu\text{l/ml}$ for the 4th and 1st instar larvae, respectively. The present studies therefore, demonstrate that moringa seed oil exhibits significant toxic action against all larval instars of Ae. aegypti especially the 1st instar. Higher levels of toxicity with increasing larval mortality were achieved with increasing dosage levels and longer exposure time.

Keywords: *Moringa oleifera* oil, *Aedes aegypti*, larval control

1. Introduction

Mosquitoes are vectors of many human diseases. According to Clements (1992) and Reinert, (2000), there are over 3000 different species of mosquitoes throughout the world. About 1,900 species occur in the humid tropics and subtropics, where the climatic conditions are favourable for rapid immature stages development and adult survival. Global efforts to reduce the number of mosquitoes usually are due to the deadly diseases they transmit to man and animals. The most important mosquito-transmitted

diseases with high public health concern include malaria, yellow fever, Dengue haemorrhagic fever and filariasis, which are transmitted by various species of Anopheles, Aedes and Culex mosquitoes.

Attempts to control these vectors have relied heavily on the use of synthetic insecticides resulting in several problems, including environmental pollution, insect resistance, unacceptable levels of residues, escalating cost of production among others (Otto, 1992). Hence, the use of plant extracts in public health and agro-ecosystem is now emerging as one of the prime methods of protecting the environment from residual toxic synthetic pesticides pollution (Prakash and Rao, 1997; Schmutterer, 1995). However, plants are by far the most efficient 'factory' for synthesis of chemical compounds, many of which they use in the defense against herbivores (Schoonhoven et al., 1998). The insecticidal metabolites from plants could therefore, be applied to other flora and fauna in the environment to protect them from injurious species.

For example, Amonkar and Reeves (1970) reported the action of garlic oil against the larval stages of five mosquito species. Badruddoza and Rahman (2008) reported the larvicidal action of moringa roots and other plants against *Aedes albopictus* and *Culex quinquefasciatus*, while Bassole et al. (2003), reported the ovicidal and larvicidal activities of leaves of three plants naturally growing in Burkina Faso. Botanical pesticides derived from members of Meliaceae family, particularly the Indian neem, *Azadirachta indica* A. Juss, are equally promising (Akou-Edi, 1984, Niber, 1994). The main active component of the neem extract is Azadirachtin, which has both deterrent, antifeedant, anti-ovipositional, growth regulatory and fecundity – reducing effects on various insects, including mosquitoes (Schmutterer, 1990). Betra et al. (1998) reported larvicidal effects of neem oil-water emulsion against *Anopheles stephensi*, *Culex quinquefasciatus* and *Ae. aegypti* in mosquito breeding habitats.

The seasonality of many plant products such as neem seeds makes it necessary to investigate as many other plants as possible. Thus many researchers are now investigating *Moringa oleifera* Lam (Horse radish) for use as pesticide. For instance, Donli and Dauda (2002) reported the use of aqueous *Moringa* seed extracts as seed dressing bio-fungicide for groundnut production in Nigeria. Adendonon et al. (2006) combined a biocontrol with *M. oleifera* seed extract for integrated control of *Sclerotium damping-off* of cowpea stem and root. Ajayi (2007) compared efficacy of *M. oleifera* seed oil with *Sesamum indicum* and *Olea europaea* against *T. castaneum*. Aqueous extracts of leaves of *M. oleifera*, *Vernonia amygdalina* and *Annona muricata* were evaluated for the control of *Collectotrichum destructivum* on seeds of cowpea (*Vigna unguiculata*) (Akinbode and Ikotun, 2008). Badruddoza and Rahman (2008) reported the larvicidal action of *M. oleifera* root extract together with other Indian plants against *Ae. albopictus* and *C. quinquefasciatus*.

The Horse radish, *M. oleifera* Lam. is an abundant invasive plant found growing throughout Nigeria and many other African countries. The many attributes of this plant have generated a lot of interests among Scientists in many African countries both for its medicinal and other uses (Ozumba, 2005). Thus its potentials as bio-pesticide need to be evaluated in different environments.

Hence, the present studies were aimed at evaluating the larvicidal action of the *M. oleifera* seed oil extract for larval control of *Ae. aegypti* mosquito.

2. Materials and Methods

2.1. Study Area

These investigations were carried out in the Research Laboratory of the Department of Parasitology and Entomology, Nnamdi Azikiwe University, Awka, Nigeria.

2.2. Rearing of Mosquito Larvae

Eggs of *Aedes aegypti* were obtained from the Federal Ministry of Health, Department of Public Health, National Arbovirus and Vectors Research Centre, Enugu, Nigeria. They were hatched and reared in the laboratory at $28 \pm 2^{\circ}\text{C}$ and $80 \pm 5\%$ r.h. The eggs were placed in transparent plastic containers containing 500mls of distilled water and allowed to hatch into 1st instar larvae and further kept to reach the 4th instar. The larvae were fed on crumbs of Yale Fortune Cabin Sweetened Biscuit; the feed was supplied every other day for normal development of the larvae.

2.3. Moringa Seed Collection and Oil Extraction

The *M. oleifera* seeds were collected from a Moringa Plantation in Enugu, Nigeria. The seeds were dried in a shed for 7 days and dehusked to obtain the kernels which were subsequently pulverized in an electric blender to obtain fine powder. Thereafter, the oil extraction was carried out using petroleum ether (Analar grade) in Soxhlet Extractor by refluxing for 3 hours. About 40g of the pulverized sample was wrapped in a double layer of Whatman No. 1 filter paper and placed inside the Soxhlet thimble. Measured volume of 250 ml of petroleum ether was poured through a funnel, by-passing the thimble containing the sample, into the round bottom flask component of the Soxhlet apparatus. This was positioned about 5 cm above an electric-heated plate while cold water was allowed to reflux in the Extractor to cool the condenser compartment. After many refluxes, the mixture of oil and ether extract was allowed to evaporate at about 75°C . The process was repeated several times at the end of which, the ether solvent distilled off and finally leaving the light yellowish oil.

2.4. Preparation of the Oil Extracts

Graded concentrations of the *M. oleifera* seed oil were prepared in acetone (Analar grade). The pure oil extract was regarded as 100% concentrate and subsequently diluted serially to give 20, 10, 5, 2.5, 0.625 and 0.3125% w/v in acetone.

2.5. Bioassay Tests

Aliquots of 1 ml of each of the above dilutions were separately placed in plastic cups containing 160 ml distilled water. Each replicate container was challenged with batches of 20 active fourth or first instar larvae of *Ae. aegypti* accordingly. Controls without oil extract were also included.

Each treatment or control was replicated five times. The bioassay tests were repeated two times at ambient laboratory temperature of $28 \pm 2^{\circ}\text{C}$, $80 \pm 5\%$ relative humidity and photoperiod 12:12 light and dark hours. Mortality / inhibition of emergence counts were made at 3 – hourly intervals for 12 hours. Dead and moribund larvae, that is, those unable to wriggle after gentle prodding were counted as dead and recorded. However, exuviae or shed cuticle started appearing in the control treatments about 2 hours post treatment but none matured into pupae or adult.

3. Data Analysis

Mortality data obtained were corrected using Abbot Formula (1925). Log_{10} versus probit regression analysis was carried out (Finney, 1972) for determining LD_{50} or level of inhibition of emergence (IE_{50}). Analysis of variance (ANOVA) was further carried out on the mortality data and means separated using least significant difference (LSD) for the different time intervals and concentration levels.

4. Results

The mortality values of 4th instar larvae of *Ae. aegypti* exposed to *M. oleifera* seed oil extract at 3 hourly intervals are presented in Table 1. The results show that moringa oil exhibited significant ($P < 0.05$) levels of toxicity to *Ae. aegypti* larvae. Mortality increased with increasing concentrations of the oil and also with increasing period of exposure. However, there were much greater responses at higher concentrations of the oil while no mortality was observed in the control treatments. Generally, dose-related mortality responses were noted in all treatments with higher ranges of the oil resulting in about 90% mortality or inhibition of emergence in the larvae, while at lower concentrations, larval mortality was low.

Dose (μl)	Mean of Mortality (%)			
	3 (hrs)	6 (hrs)	9 (hrs)	12 (hrs)
200	5	20	20	90
100	5	10	55	85
50	10	20	55	80
25	5	15	35	65
12.5	5	15	30	60
6.25	5	10	20	45
3.125	5	10	15	25
Mean	$5.7 \pm 1.2\text{a}$	$14.3 \pm 2.2\text{a}$	$32.1 \pm 6.5\text{b}$	$65.0 \pm 7.0\text{c}$
Control	0	0	0	0

Table 1: Percentage mortality of 4th instar larvae of *Aedes aegypti* exposed to moringa seed oil at various time intervals.

LSD = 12.0

*Mean of five replicates (\pm s.e)

*The means followed by the same letter in column are not significantly different from each other ($P < 0.05$) by LSD.

The probit regression analysis (Figure 1) showed that after 12 hours of exposure, the LD_{50} (IE_{50}) of the moringa oil – treated 4th instar larvae was $208.35 \mu\text{l}$ and $62.84 \mu\text{l}$ for 1st instar larvae.

Table 2 also shows that mortality of 1st instar larvae of the mosquito increased significantly ($P < 0.05$) with increasing levels of oil concentration and also with increasing period of exposure, giving an LD_{50} (IE_{50}) value of the oil for this instar as $62.94 \mu\text{l}$.

Mortality of both 4th and 1st instar larvae (Table 1) after the 6th hour was significantly ($P < 0.05$) higher than at the first 3 – hours. This trend was observed in all the treatments irrespective of the dosage applied. However, at the highest concentration of $200 \mu\text{l}$, the mean toxicity of the oil to the 4th instar was 49% while that of the 1st instar was 66.5% (Table 3). The LD_{50} values determined at the 3 hourly intervals are shown in Figure 2. This indicates that exposure of the 4th instar larvae for 3, 6, 9 and 12 hours would require 2.29×10^{38} (i.e. α), 2.76×10^7 , 4.08×10^2 and $1.35 \times 10^1 \mu\text{l}$ of moringa oil respectively, order to achieve the same level of 50 percent kill. Similarly, the 1st instar larvae would require exposure to 3.85×10^3 , 7.11×10^1 , 3.45×10^1 and $1.22 \times 10^1 \mu\text{l}$ of the oil extract for 3, 6, 9 and 12 hrs, respectively, to kill 50 percent of the larvae (Figure 3). These are all indications that the early instars are more susceptible and require less toxicant to kill than later instars, irrespective of the duration of exposure.




Figure 1: Toxicity of Moringa seed oil to 4th and 1st instar larvae of Aedes aegypti

- ◆ 4TH INSTAR.....y = 0.5558 x + 3.7112
R² = 0.9517, LD₅₀ = 208.35 µl
- 1ST INSTAR.....y = 0.6311 x + 3.8647
R² = 0.9727, LD₅₀ = 62.94 µl.

Dose (µl)	Mean of Mortality (%)			
	3 (hrs)	6 (hrs)	9 (hrs)	12 (hrs)
200	25	70	80	100
100	25	50	60	75
50	30	40	45	65
25	25	35	40	50
12.5	25	30	40	45
6.25	10	25	30	40
3.125	10	20	25	30
Mean	21.43±1.5a	38.57±2.5b	45.71±5.4c	57.9±7.0d
Control	0	0	0	0

Table 2: Percentage mortality of 1st instar larvae of Aedes aegypti exposed to moringa seed oil at various time intervals.

LSD = 8.66

*Mean of five replicates (± s.e)

*The means followed by the same letter in column are not significantly different from each other (P<0.05) by LSD.

However, lower concentrations for longer periods of contact with the larvae would achieve similar level of kill compared with higher concentrations applied for shorter periods of time.

Dose (µl)	Mean of Mortality (%)	
	4 th Instar	1 st Instar
200	49.0	66.5
100	38.8	52.5
50	41.3	45.0
25	30.0	37.5
12.5	27.5	35.0
6.25	20.0	26.3
3.125	13.8	21.3
Control	0	0
Mean (±s.e.)	28.63 ± 5.6a	40.59 ± 5.4 b
LSD = 8.64		

Table 3: Cummulative Toxicity of Moringa seed oil to 4th and 1st instar larvae of Aedes aegypti, after 12 hours

Mean of five replicates (\pm s.e).

Means followed by the same letter(s) in column are not significantly different from each other ($P < 0.05$).




Figure 2: Mortality of 4th instar larvae of *Aedes aegypti* exposed to moringa seed oil at various time intervals

- ◆ **3 HOURS** $y = 0.0431x + 3.3467$
 $R^2 = 0.0417$, $LD_{50} = (2.29 \times 10^{38}) \mu l$
- **6 HOURS**..... $y = 0.1797x + 3.6629$
 $R^2 = 0.3427$, $LD_{50} = (2.76 \times 10^7) \mu l$
- ▲ **9 HOURS**..... $y = 0.376x + 3.9918$
 $R^2 = 0.2726$, $LD_{50} = (4.08 \times 10^2) \mu l$
- × **12 HOURS**..... $y = 0.8785x + 4.0066$
 $R^2 = 0.5942$, $LD_{50} = (1.35 \times 10^1) \mu l$




Figure 3: Mortality of 1st instar larvae of *Aedes aegypti* exposed to moringa seed oil at various time intervals

- ◆ **3 HOURS**..... $y = 0.3779 x + 3.6451$
 $R^2 = 0.6062, LD_{50} = (3.85 \times 10^3) \mu\text{l}$
- **6 HOURS**..... $y = 0.6784 x + 3.7438$
 $R^2 = 0.9249, LD_{50} = (7.11 \times 10^1) \mu\text{l}$
- ▲ **9 HOURS**..... $y = 0.7416 x + 3.8595$
 $R^2 = 0.8854, LD_{50} = (3.45 \times 10^1) \mu\text{l}$
- × **12 HOURS**..... $y = 1.8549 x + 2.9863$
 $R^2 = 0.6324, LD_{50} = (1.22 \times 10^1) \mu\text{l}$

5. Discussion

The larvicidal toxicity of *M. oleifera* seed oil to both 1st and 4th instar larvae of *Ae. aegypti* was studied in the laboratory. The studies have shown that this insecticide of plant origin can be used as independent control method in reducing the impact of mosquitoes. Considering the high cost of conventional insecticides and other environmental concerns, moringa seed oil could provide a cheap alternative means of mosquito vector control. Moringa has added advantage of being comparatively harmless to man and other non-target organisms. It could, therefore, with little education, be cultivated and applied to major vector breeding sites by communities.

In the present studies, the general performance of moringa oil in killing the larvae of *Ae. aegypti* was observed with relatively high percentage larval mortality recorded in both 1st and 4th instar larvae, high percentage inhibition of emergence and LD_{50} values. The toxic effect of the oil was high, as the mortality and the rate of inhibition of emergence increased with increasing concentration and time. The effectiveness of plant oils in regulating growth and inhibition of many insects and fungal development has been reported previously (Jilani and Su, 1983, Mulla et al., 2003). For instance, Manas et al. (2005) reported that *M. oleifera* extracts inhibited 97.32% germination of fungal pathogen. Sharma and Saxena (1994) also found that the petroleum ether extract of *Tagetes erectes* had toxic effect on larvae of *Anopheles stephensi* and on its significant growth index. Mwangi and Mukiyama (1988) observed that fraction of *Melia volkensii* fruit kernel extract had growth inhibition activity at low concentration, whereas two other fractions had acute toxic effects on the mosquito larvae. Thus, in the present studies only a few surviving larvae were left in some treated samples, and they were unable to mature into the next stage of development. Pushpalatha and Muthukrishnam (1995) reported that leaf extracts of *Vitex negundo* at very low concentrations had larvicidal activity against *Culex quiquefasciatus* and *A. stephensi* and also extended the duration of larval pupation.

The botanical pesticide, Moringa seed oil, has proved to be very effective in mosquito larval control as demonstrated in the present studies. The activity tended to extend over a longer period, and its effectiveness increased with increasing exposure time. The oil probably exhibited its effects on the mosquito larvae as growth regulator, as well as blocking the respiratory organs. According to Hirashima et al. (1998) and Enan (2001) some plant essential oils block the octopamine neuroreceptors that regulate the movement, heart rate, behavior, metabolism, and pupation of insects. Thus, the regulatory activity of Moringa seed oil was exhibited in the mortality of the larvae and inhibition of their eclosion into pupal stage.

The present studies have further demonstrated that the 1st instar larvae of *Ae. aegypti* were more susceptible than the 4th instar. This observation agrees with reports that early instars of insects are more susceptible to inhibitory and growth regulatory effects of most insecticides (Mulla et al., 2003). This, the authors however, suggested was related to the size and tender nature of the chitin in the early instars of the mosquitoes. Furthermore, Akpa et al. (2003) observed that 3rd instar was more susceptible than 4th instar when a plant oil of *Phytolacca dodecandra* was used against *Ae. aegypti*.

The present studies therefore, demonstrate that *M. oleifera* could be used as mosquito larvicide for the control of *Ae. aegypti*. This would help to contain any outbreak of the yellow fever often associated with the mosquito (Lee and Moore, 1972, Einterz, 1971). It would therefore, be necessary to increase campaign on the growing of the crop in all ecological zones of Nigeria, as an invaluable tool for the control of mosquito vectors and hence the diseases they transmit.

6. References

1. Abbot, W.A. (1925). A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology*, 18:265-267.
2. Adendonon, A., Aveling, T.A.S., Labusehanone, N. and Tamo, M. (2006). Biocontrol agent in combination with *Moringa oleifera* Extract for Integrated Control of *Sclerotium* – Cause of Cowpea Damping off and stem rot. *European Journal of Plant Pathology*, 115(4): 409-418.
3. Ajayi, F.A. (2007). Effect of grain breakage and application of edible oils to protect pearl millet, *Pennisetum glaucum* (L.) R. Br., against infestation by adult *Tribolium castaneum* (Herbst.) (Coleoptera: Tenebrionidae) in Lafia, Nigeria. *Nig. J. Ent.*, 24:107-113.
4. Akinbode, O.A. and Ikotun, T. (2008). Efficacy of certain plant extracts against seed-borne infection of *Collectotrichum destructivum* on cowpea (*Vigna unguiculata*). *African Journal of Biotechnology*, 7(20): 3683-3685.

5. Akou-edi, D. (1984). Effects of neem seed powder and oil on *Tribolium confusum* and *Sitophilus zeamais*. In: Schumutterer, H., Ascher, K.R.S. (eds). Natural Pesticides from the neem tree (*Azadirachta indica* (A.) (Juss) and other tropical plants. Proceedings of the Second International Neem conference, Rauischolzhausen, Federal Republic, Deutsche Gexllanchoff Fir, Technishe Zucammena Bat, Pp. 445-452.
6. Akpa, A.U.C., Mwale, N.A., Ozumba, N.A. and Yohannes, L.W. (2003). The efficacy of *Phytolacca dodecandra* (L.) (Herit) as a larvicide against *Aedes aegypti* (Linnaeus) – a yellow fever vector in Nigeria. *Journal of Science and Technology*, 9:13-22.
7. Amonkar, S.V. and Reeves, E.L. (1970). Mosquito control with active principle of garlic, *Allium sativum*. *Journal of Economic Entomology*, 63: 1172 – 1175.
8. Badruddoza. D.M. and Rahman, S.M.K. (2008). Lavicidal activities of aqueous extracts of nineteen Indian plants against two mosquitoes, *Aedes albopictus* and *Culex quinquefasciatus*. *Crop Protection and Toxicology*, 4:19-25.
9. Bassole, I.H., Guelbeogo, W.M., Nebie, R., Constantini, R., Sagnon, N., Kabore, Z.I. and Traore, S.A. (2003). Ovicidal and larvicidal activity of essential oils extract from three spontaneous plant against *Aedes aegypti* and *An. gambiae* complex mosquitoes of Burkina Faso. *Parasitologia*, 45(1): 23-26.
10. Betra, C.P., Mital, P.K., Adak, T and Sharma, V.P. (1998). Efficacy of neem oil water emulsion against mosquito immature. *Indian Journal of Malariology*, 35(1): 15-21.
11. Clements, A. N. (1992). The biology of mosquitoes: (Vol. 1) Development, nutrition and reproduction. Chapman and Hall, New York. P126 – 135.
12. Donli P.O. and Dauda, H. (2002). Evaluation of Acqeous Moringa seed extracts as a seed treatment biofungicide for groundnuts. *Pest Management Science*. 59:9.
13. Einterz, H. (1971), “Yellow fever” in Eastern Nigeria. *The Lancet*, July 2, 39-40.
14. Enan, E. (2001). Insecticidal activity of essential oils: octopaminergic sites of action. *Comp. biochem. Physiol.*, 130:325-337.
15. Finney, D.J. (1971). Probit analysis. Ed. 3. Cambridge University Press, London.
16. Hirashima, A.S., Hirokado, R., Tojikubo, R., Takeya, E., Taniguchi, and Eto, M. (1998). Metamorphosis of the red flour beetle, *Tribolium Freeman Honton* (Coleoptera: Tenebrionidae): alteration of octopamine content modulates activity of juvenile – hormone esterase, ecdysteriod level, and pupation. *Arch. Insect Biochem. Physiol.*, 37:33 – 46.
17. Jilani, G. and Su, H.C.F. (1983). Laboratory studies on several plant materials as insect repellents for protection of cereal grains. *J. Econ. Entomol.*, 76: 154 – 157.
18. Lee, V.H. and Moore, D.L. (1972). Vectors of the 1969 Yellow Fever epidemic on the Jos Plateau, Nigeria. *Bulletin of the World Health Organization*, 46:669-673.
19. Manas, D.M., Souman, C., Pratheesh, K., Beera, S. (2005). In vitro screening of some plant extract against fungal pathogens of mulberry (*Morus* spp.). *Archives of Phytopathology and Plant Protection*, 38(3):157-164.
20. Mulla, M.S., Thavura, U., Tawatsin, A., Chompoonsri, J., Zaim, M. and Su, T. (2003). Laboratory and Field evaluation of Novaluron, a new acylurea insect growth regulator; against *Aedes aegypti* (Diptera: Culicidae). *J. Vector Ecol.*, 28:241-254.
21. Mwangi, R.W. and Mukhiana, T.K. (1988). Evaluation of *Melia volkensi* extract fractions as mosquito larvicides. *J. Amer. Mosq. Cont. Assoc.*,4:442-447.
22. Niber, B.T. (1994). The ability of powders and slurries from ten plant species to protect stored grain from attack by *Prostepahnus truncatus* (Horn.) (Coleoptera: Bostrichidae) and *Sitophilus oryzae* L. (Coleoptera: Curculionidae). *J. Stored prod. Res.*, 30:297-301.
23. Otto, D. (1992). Insektizidresisyenz in Insektenpopulation – en-ein uberblick. *Nachrichteblatt des Dt. Pflanzen – Schutzdienstes*, 44(2)34-39. (Abstract only).
24. Ozumba, N.A. (2005). “Health and Nutrition” An invited paper presented at a workshop on skills Enhancement for Traditional Medicine Practitioners. Organized by Nigeria National Medicine Development Agency, Federal Ministry of Science and Technology. Modotel, Enugu, 24-26 January. 2005.
25. Prakash, A. and Rao, J. (1997). Botanical Pesticides in Agriculture. CRC Lewis publishers Boca Raton. 150 pp.
26. Pushpalatha, E. and Muthukrishnan, J. (1995). Larvicidal activity of few plant extracts against *Culex quinquefasciatus* and *Anopheles stephensi*. *Ind. J. Malariol.*, 32:14-23.
27. Reinert, J.F. (2000). New classification for the composite genus *Aedes* (Diptera: Culicidae: Aedini) elevation of the subgenus *Ochleratatus* to generic rank, reclassification of the other subgenera, and notes on certain subgenera and subspecies. *Journal of American mosquito Control Association*, 16:175-188.
28. Schmutterer, H. (1995). The neem tree: source of unique natural products for the integrated pest management, medicine, industry and other purposes. Vch Verlagsgesellschaft, Weinheim.
29. Schmutterer, H. (1990). Properties and potentials of natural pesticides from the neem tree, *Azadirachta indica*. *Annu. Rev. Entomol.*, 35:271-297.
30. Schoonhoven, L.M.; Jermy, T. and Van Loon, J.A.A. (1998). *Insect Plant Biology: From physiology to evolution*: Chapman & Hall, London.
31. Sharma, M. and Saxena, R.C. (1994). Phytotoxicological activity of *Tagetes erectes* in aquatic stages of *Anopheles stephensi*. *Ind. J. Malariol.*, 31:21-26.